



Standard Test Method for Total Chlorine in Epoxy Resins and Compounds¹

This standard is issued under the fixed designation D4301; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This test method describes a procedure for the determination of total chlorine in epoxy resins and glycidyl ethers.

1.2 The values stated in SI units are to be regarded as standard. No other units of measurement are included in this standard.

1.3 *This standard does not purport to address all of the safety concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use. For specific hazard statements, see Section 8.*

2. Referenced Documents

2.1 *ASTM Standards:*²

D1193 Specification for Reagent Water

D6440 Terminology Relating to Hydrocarbon Resins

E691 Practice for Conducting an Interlaboratory Study to Determine the Precision of a Test Method

2.2 *Other Document:*

OSHA Regulations, 29 CFR paragraphs 1910.1000 and 1910.1200³

3. Terminology

3.1 *Definitions:* For definitions of terms used in this standard, See Terminology D6440.

4. Summary of Test Method

4.1 The material, dissolved in dimethoxyethane (DME) or other suitable inert solvent, is reacted with sodium biphenyl to convert bound organic chlorine to the water soluble chloride.

¹ This test method is under the jurisdiction of ASTM Committee D01 on Paint and Related Coatings, Materials, and Applications and is the direct responsibility of Subcommittee D01.33 on Polymers and Resins.

Current edition approved Nov. 1, 2013. Published November 2013. Originally approved in 1984. Last previous edition approved in 2005 as D4301 – 05. DOI: 10.1520/D4301-05R13.

² For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For *Annual Book of ASTM Standards* volume information, refer to the standard's Document Summary page on the ASTM website.

³ A suitable mechanical holder is available from the Gardner Laboratory, Inc., 5521 Landy Lane, Washington, DC, Item 660.

The excess reagent is decomposed with isopropyl alcohol. The chloride ion is then titrated potentiometrically with silver nitrate.

5. Significance and Use

5.1 The presence of residual chlorine in epoxy resins is deleterious to final product properties. This test method has been found to be applicable to resins or ethers with chlorine contents ranging from 50 ppm to 35 % by weight. Other halogen compounds react with the reagent but are distinguished from chlorine by the final potentiometric titration. Epoxy and other functional groups will consume reagent but do not affect the results.

6. Apparatus

6.1 *Potentiograph.*

6.2 *Beaker*, of appropriate size.

6.3 *Buret*, of appropriate size.

6.4 *Stirrer*, magnetic or paddle.

7. Reagents

7.1 *Purity of Reagents*—Reagent grade chemicals shall be used in all tests. Unless otherwise indicated, it is intended that all reagents shall conform to the specifications of the Committee on Analytical Reagents of the American Chemical Society, where such specifications are available.⁴ Other grades may be used, provided it is first ascertained that the reagent is of sufficiently high purity to permit its use without lessening the accuracy of the determination.

7.2 *Purity of Water*—Unless otherwise indicated, references to water shall be understood to mean reagent water conforming to Type II of Specification D1193.

⁴ *Reagent Chemicals, American Chemical Society Specifications*, American Chemical Society, Washington, DC. For suggestions on the testing of reagents not listed by the American Chemical Society, see *Analar Standards for Laboratory Chemicals*, BDH Ltd., Poole, Dorset, U.K., and the *United States Pharmacopeia and National Formulary*, U.S. Pharmacopeial Convention, Inc. (USPC), Rockville, MD.